Product Nume         HurriPAK" Proiodontal Anesthetic Stater KI         Cetacaine® Topical Anesthetic Liquid         Cetacaine® Topical Anesthetic Liquid         Constitution (Liquid)         Constethisticon (Liquid)         Constethisticon (Liquid) <th>Nee</th> <th>dle-free Periodon</th> <th>tal Anesthesia Co</th> <th>mparison Chart</th>	Nee	dle-free Periodon	tal Anesthesia Co	mparison Chart
Packaging Packaging         Included in kit - 2 bottle of fund - 2 bottle of fund - 2 digraphic burgets - 1 digraphic burgets - 2 digraphic burg		HurriPAK <sup>™</sup> Periodontal Anesthetic	Cetacaine® Topical Anesthtetic Liquid	
Packaging <ul> <li>2 bit field of thursdame Topical Anesthetic Light of thursdame Linguist of thursdame Linguist Instance <ul> <li>2 displace to the Linguist</li></ul></li></ul>	Manufacturer	Beutlich <sup>®</sup> Pharmaceuticals, LLC	Cetylite <sup>®</sup> Industries, Inc.	Dentsply Pharmaceutical
Approximate Coal         Complete kit: \$109.95         Complete kit: \$199.99         et also participation (************************************	Packaging	<ul> <li>2 bottles of HurriCaine® Topical Anesthetic Liquid (1 fl. oz./29.6 mL) Original Wild Cherry and Piña Colada flavors</li> <li>12 disposable irrigation syringes</li> <li>12 disposable plastic periodontal irrigation tips</li> </ul>	<ul> <li>30 g bottle of liquid</li> <li>1 Luer-lock cap</li> <li>100 Microcapillary Delivery Tips</li> </ul>	<ul> <li>20 cartridges containing</li> <li>1.7 g gel and tips</li> <li>Unique dispenser - autoclaving recommended</li> </ul>
Approximate Cost         Complete kit: \$199.99         • Ocarridges and tips: \$11.99           Approximate Packaging         • HurriCaine Topical Anstetic Liquid - 1 a. co. totik: will difference of the codars: \$9.40         • 100 carridges and tips: \$11.99           Approximate Retifing         • HurriCaine Topical Anstetic Liquid - 1 a. co. totik: will difference of the codars: \$9.40         • 20 carridges and tips: \$11.99           Approximate Retifing         • HurriCaine Topical Anstetic Liquid - Status         • 100 Microcapilary Delivery Tips: \$99.00         20 carridges and tips: \$11.99           Active Ingredencity         20% bancocaline         14% Benzocaline, 2% Tescale hydrochionic         2.5% Lidocaline and 2.5% Prilocaline           Systemic         Virtually none         Yes         2.5% Lidocaline and 2.5% Prilocaline           OTC or Pk         OTC         Pk         Rk           Contraindications         Contrained participant to approximate to appendix to append	Flavor(s)	Original Wild Cherry AND Piña Colada Flavors	Banana	Unflavored
Approximate Original Packaging         11.8.c. bottle, Wild Cherry or Prinz Colada:: 93.6.6 Hurri2AR etails in patient patient inglation syntagis & plastic periodontal imgation tips - 20 tips - 30.3 bottle of liquid:: \$35.00         20 cartridges and tips: \$114.99           Packaging         + Hurri2AR etails - H		Complete kit: \$109.95	Complete kit: \$199.99	<ul> <li>20 cartridges and tips: \$114.99</li> </ul>
Ingredient(s)         20% Benzobarine         Tetracaine Hydrochloride         2.3% Lubocalite and 2.3% Prilocalite           Systemic Absorption         Virtually none         Yes         Yes         Rx           OTC or Rx         OTC         Rx         Rx         Rx           Contraindications         Catacaline is not suitable and should never be used for injection. Do not use on the eyes. To avoid excessive systemic absorption, Cetacaline should not be applied to a graphen should not be administered to patients with a structure of the product.         Patients with a known history of hypersensitivity to local anesthetics of the amide type or to any other component of the product.           Contraindications         Methemoglobinemia has been reported following the use of benzocaline on extremely rare occasions. Intravenous enthylene blue is the specific therapy for this condition.         Patients with glucose-6-phosphate dehydrogenase deficiency or congenital or idiopatitic methemoglobinemia are more should be observed when using Catacaline should be observed when using Catacaline the specific therapy for this condition.         Patients with glucose-6-phosphate dehydrogenase deficiency or congenital or idiopatitic methemoglobinemia are or complete information.           Precuutions         On rare occasions, methemoglobinemia has been reported in connection with the use of other connection with the use of obsercoalme-containing products.         Orapix should be used with caution in commended dosage (see Dosage and Administration). If a patient becomes construct the specific therapy for this condition.         Orapix should be used with caution in commended dosage (see Dosage and Adminis	Cost of Refill	<ul> <li>1 fl. oz. bottle, Wild Cherry or Piña Colada: \$9.46</li> <li>HurriPAK Refill Kit - 10 disposable irrigation syringes &amp; plastic periodontal irrigation tips: \$26.45</li> <li>HurriPAK Periodontal Irrigation Tips - 20 tips: \$33.45</li> </ul>	100 Microcapillary Delivery Tips: \$99.00	20 cartridges and tips: \$114.99
Absorption         Virtually noise         Yes         Tes           OTC or Rx         OTC         Rx         Rx           Cottor Rx         OTC         Rx         Rx           Cottor Rx         OTC         Rx         Rx           Cottor Rx         OTC         Rx         Rx           Contraindications         Editable and should not be applied to large areas of demunded administered to patient shound not be applied to large areas of demunded administered to patient shound not be applied to large areas of demunded administered to patient shound not be applied to patient shound not be applied to large areas of demunded to be administered to patient shound not be applied to large areas of demunded to be administered to patient shound not be applied to applied the domunded to be administered to patient shound not be applied to applied the domunded to be administered to patient shound not be applied to applied the domunded to be administered to applied the domunded to app		20% Benzocaine		2.5% Lidocaine and 2.5% Prilocaine
OTC or Rx         OTC         Rx         Rx           OTC or Rx         OTC         Rx         Rx           Contraindications         Cetacaine in ont suitable and should never be used for injection. Do not use on the eyes. To avoid excessive systemic absorption, Cetacaine should not be applied to large areas of denuded or inflamed tissue. Cetacaine should not be applied to large areas of denuded or inflamed tissue. Cetacaine should not be to administered to galaties the applied to the applied		Virtually none	Yes	Yes
Contraindications         Celetacine is not uibble and should never be used on research. Do not use only in the granted to the applied to large areas of cliculate should not be applied to large areas of cliculate administered to patients who are hypersensitive to any of its ingredients or to patients how on to have cholinestered to patients who are hypersensitive to administered to patients who are receiving treatments with methemoglobinemia are of barocacine-containing products. Care been reported in connection with the use of berzocaine-containing products. Care been reported in connection with the use of barocacine-containing products. Care been reported in connection with the use of barocacine-containing products. Care been reported in connection with the use of barocacine-containing products. Care been reported in connection with the use of barocacine-containing products. Care been reported in connection with the use of barocacine-containing products. Care been reported in connection with the use of barocacine-containing products. Care been reported in connection with methaling products. Care been reported in connection with		OTC	Bx	Bx
Warnings       Methemoglobinemia has been reported following the use of benzocaine on extremely rare occasions. Intravenous methylene blue is the specific therapy for this condition.       Image: Condition of the transmission of transmission of transmission of transmission of the transmission of transmit transmissin of the transmission of transmit transmission of tra	Contraindications		used for injection. Do not use on the eyes. To avoid excessive systemic absorption, Cetacaine should not be applied to large areas of denuded or inflamed tissue. Cetacaine should not be administered to patients who are hypersensitive to any of its ingredients or to patients known to have cholinesterase deficiencies. Tolerance may vary with the status of the patient. Cetacaine should not be used under dentures or cotton rolls, as retention of the active ingredients under a denture or cotton roll could possibly cause an escharotic effect. Routine precaution for the use of any topical anesthetic	
PrecautionsOn rare occasions, methemoglobinemia has been reported in connection with the use of benzocaine-containing products.been reported in connection with the use of of benzocaine-containing products. Care should be used not to exceed the maximum recommended dosage (see Dosage and Administration). If a patient becomes cyanotic, treat appropriately to counteract (such as with methylene blue, if medically indicated).Oradix should be used with califormia on to combination with dental injection anesthesia, other local anesthetics; e.g., Class 1 antiarrhythmics such as tocainide and mexiletine, as the toxic effects of these drugs ar likely to be additive and potentially synergistic.On setWithin 30 seconds30-60 seconds30 secondsDuration15 minutes30-60 minutes20 minutesStorage20-25 °C (68-77 °F)Store in a cool dry place25 °C (77 °F) – If it has turned to gel, it should be refrigerated until it becomes a liquid.Irrigation TipsPre-angled, end port disposable plastic periodontal irrigation tips • Gentle to soft tissue • Does not look like a needle – patients accept it • Easy to maneuver to get deep within the sulcusMinimally invasive plastic tipsSterie end port, blunt, metal tips • Breakage more likely if bent at hub • Requires sharps container for disposal (per manufacturer instructions)StringeDispaceable 2 ml euripageDispaceable 0.4 ml euripagePrefilled cartridges and customized,	Warnings	following the use of benzocaine on extremely rare occasions. Intravenous methylene blue is		dehydrogenase deficiency or congenital or idiopathic methemoglobinemia are more susceptible to drug-induced methemoglobinemia. Oraqix should not be used in those patients with congenital or idiopathic methemoglobinemia and in infants under the age of twelve months who are receiving treatments with methemoglobin- inducing agents. See product labeling insert for
Duration       15 minutes       30-60 minutes       20 minutes         Storage       20-25 °C (68-77 °F)       Store in a cool dry place       25 °C (77 °F) – If it has turned to gel, it should be refrigerated until it becomes a liquid.         Irrigation Tips       Pre-angled, end port disposable plastic periodontal irrigation tips       Store in a cool dry place       Sterile end port, blunt, metal tips         Irrigation Tips       Ones not look like a needle – patients accept it easy to maneuver to get deep within the sulcus       Minimally invasive plastic tips       Sterile end port, blunt, metal tips         Stringer/Contrided       Disposable 2 mL suringer       Disposable 2 mL suringer       Prefilled cartridges and customized,	Precautions	been reported in connection with the use of	been reported in connection with the use of benzocaine-containing products. Care should be used not to exceed the maximum recommended dosage (see Dosage and Administration). If a patient becomes cyanotic, treat appropriately to counteract (such as with	combination with dental injection anesthesia, other local anesthetics, or agents structurally related to local anesthetics; e.g., Class 1 antiarrhythmics such as tocainide and mexiletine, as the toxic effects of these drugs are
Storage       20-25 °C (68-77 °F)       Store in a cool dry place       25 °C (77 °F) – If it has turned to gel, it should be refrigerated until it becomes a liquid.         Irrigation Tips       Pre-angled, end port disposable plastic periodontal irrigation tips       Store in a cool dry place       25 °C (77 °F) – If it has turned to gel, it should be refrigerated until it becomes a liquid.         Irrigation Tips       Pre-angled, end port disposable plastic periodontal irrigation tips       Minimally invasive plastic tips       Sterile end port, blunt, metal tips         • Gentle to soft tissue       • Does not look like a needle – patients accept it       Minimally invasive plastic tips       Sterile end port, blunt, metal tips         • Requires sharps container for disposal (per manufacturer instructions)       • Disposable 0.4 mL suringes       Prefilled cartridges and customized,	Onset	Within 30 seconds	30-60 seconds	30 seconds
Storage       20-25 C (65-77 P)       Store in a cool of y place       refrigerated until it becomes a liquid.         Irrigation Tips       Pre-angled, end port disposable plastic periodontal irrigation tips       Store in a cool of y place       Sterile end port, blunt, metal tips         Irrigation Tips       Gentle to soft tissue       Minimally invasive plastic tips       Sterile end port, blunt, metal tips         • Does not look like a needle – patients accept it       • Disposable 0.4 mL suringes       Prefilled cartridges and customized,	Duration	15 minutes	30-60 minutes	
Irrigation Tips       periodontal irrigation tips       Minimally invasive plastic tips       Sterile end port, bluit, metal tips         Irrigation Tips       Gentle to soft tissue       Minimally invasive plastic tips       Breakage more likely if bent at hub         • Does not look like a needle – patients accept it       Minimally invasive plastic tips       Prequires sharps container for disposal (per manufacturer instructions)         Suringe/Cortridge       Disposable 3 mL suringes       Prefilled cartridges and customized,	Storage	20-25 °C (68-77 °F)	Store in a cool dry place	25 °C (77 °F) – If it has turned to gel, it should be refrigerated until it becomes a liquid.
	Irrigation Tips	periodontal irrigation tips • Gentle to soft tissue • Does not look like a needle – patients accept it	Minimally invasive plastic tips	<ul> <li>Breakage more likely if bent at hub</li> <li>Requires sharps container for disposal</li> </ul>
	Syringe/Cartridge	Disposable 3 mL syringes	Disposable 0.4 mL syringes	



HurriCaine is a registered trademark of Beutlich Pharmaceuticals, LLC. HurriPAK is a trademark of Beutlich Pharmaceuticals, LLC. Cetacaine is a registered trademark of Cetylite Industries, Inc. Oraqix is a registered trademark of Dentsply Pharmaceutical. HPCC 430 1214

## **STARTER KIT**

GREAT TASTING ORIGINAL WILD CHERRY AND PIÑA COLADA FLAVORS!



## HURRIPAK<sup>™</sup> PERIODONTAL ANESTHETIC STARTER KIT

Only use what you need - no pre-filled carpules No expensive dispensers or accessories to buy

HurriPAK<sup>™</sup> Periodontal Anesthetic Kit is a great alternative to local injections prior to root planing and scaling or full mouth debridement. Whether scaling an entire quadrant or an isolated area, HurriPAK allows dispensing only the amount of liquid needed, so no product is wasted. Plastic irrigation tips enable effective sub-gingival application of HurriCaine<sup>®</sup> Topical Anesthetic Liquid and are gentle to soft tissue.

FEATURES	BENEFITS	
Fast onset	Works within 30 seconds	
OTC product	Safer than Rx products	
Plastic irrigation tips	Accepted well by patients	
Disposable syringes and tips	Convenient	
Original Wild Cherry and Piña Colada flavors	Pleasant tasting	
More economical than other kits	Saves you money!	

## **ORDERING INFORMATION:**

0283-1009-09 HurriPAK Periodontal Anesthetic Starter Kit Contains: 1 fl. oz. bottle of Wild Cherry flavored HurriCaine Liquid, 1 fl. oz. bottle of Piña Colada flavored HurriCaine Liquid, 12 disposable periodontal irrigation syringes and tips and 2 Luer-Lock Dispensing Caps 0283-1107-07 HurriPAK Refill Kit Contains: 10 disposable periodontal irrigation syringes and tips

0283-0908-21 HurriPAK Periodontal Irrigation Tips Package Contains: 20 disposable periodontal irrigation tips



To place an order or for more information, please contact your Dental Supplier or call us at 1-800-238-8542, M-F: 8:00 a.m. – 4:30 p.m. ET. HurriCaine is a registered trademark of Beutlich® Pharmaceuticals, LLC. HurriPAK is a trademark of Beutlich Pharmaceuticals, LLC. www.beutlich.com