

HURRIPAK™ Periodontal Anesthetic Kit

20% Benzocaine Oral Anesthetic Liquid

Drug Facts

Active ingredient

Benzocaine 20% Oral Anesthetic

Purpose

Uses for the temporary relief of occasional minor irritation and pain associated with minor dental procedures

Warnings

Allergy alert: Do not use if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

When using this product • avoid contact in eyes

Keep out of reach of children. If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center immediately.

Directions:

- do not exceed recommended dosage
- adults and children 12 years and older: pull up to 3 mL of HurriCaine liquid into the disposable syringe and then attach the disposable tip. Apply a small amount of HurriCaine liquid to the gingival margin and directly into the periodontal pockets on the buccal and lingual surfaces, one quadrant at a time. Anesthesia will take place in 15 - 30 seconds.
- 3 mL or less should be adequate for complete scaling of all 4 quadrants
- discard syringe and tip after use on each patient

Other information

- Do not use if imprinted seal under cap of HurriCaine Liquid is broken or missing
- store at 20-25 °C (68-77 °F)

Inactive ingredients flavor, polyethylene glycol, sodium saccharin

Questions or comments?

1-800-238-8542 M-F: 8:00 a.m. – 4:30 p.m. ET

Beutlich® Pharmaceuticals, LLC

7770 S. US Hwy 1, Suite H, Bunnell, FL 32110

HurriCaine® is a registered trademark of Beutlich Pharmaceuticals, LLC.

HurriPAK™ is a trademark of Beutlich Pharmaceuticals, LLC.

www.beutlich.com HPDF 467 1214

