

# HURRICAINE® Topical Anesthetic Spray

20% Benzocaine Oral Anesthetic

**1/2 Second Spray Is All You Need!**

## Drug Facts

### Active ingredient

Benzocaine 20% ..... Oral Anesthetic

### Purpose

**Uses** for the temporary relief of occasional minor irritation and pain, associated with:

- canker sores
- sore mouth and throat
- minor dental procedures
- minor irritation of the mouth and gums caused by dentures or orthodontic appliances
- minor injury of the mouth and gums

### Warnings

**Allergy alert:** Do not use if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

**Flammable. Do not use near fire or flame. Avoid smoking during use.**

### Stop use and ask a doctor if

- sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting
- sore mouth symptoms do not improve in 7 days, or if irritation, pain, or redness persists or worsens

### When using this product

- avoid spraying in eyes
- do not store at temperature above 49 °C (120 °F)
- do not puncture or incinerate. Contents under pressure.
- use only as directed. Intentional misuse by deliberately concentrating and inhaling the contents can be harmful or fatal.

**Keep out of reach of children.** If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center immediately.

**Adverse Reactions:** Methemoglobinemia has been reported following the use of benzocaine on extremely rare occasions. Intravenous methylene blue is the specific therapy for this condition.

### Directions:

- do not exceed recommended dosage
- firmly insert extension tube into spray can valve and ensure it fits securely into the actuator opening
- position the extension tube 1-2 inches from the affected area
- spray 1/2 second (press and immediately release actuator)
- repeat once if necessary
- anesthesia is accomplished in approximately 20 seconds

adults and children 2 years of age and older	use up to 4 times daily or as directed by a dentist or doctor
children under 12 years of age	should be supervised in the use of the product
children under 2 years of age	consult a dentist or doctor

### Inactive ingredients

flavor, hydrocarbon propellant (isobutane and propane), polyethylene glycol, sodium saccharin

### Questions or comments?

**1-800-238-8542** M-F: 8:00 a.m. – 4:30 p.m. ET